

## Live attenuated vaccine, avian coccidiosis

### STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER INGREDIENT(S):

#### EVANT

#### Active substances:

Each 0.007 ml dose contains:

<i>Eimeria acervulina</i> , strain 003	332 – 450*
<i>Eimeria maxima</i> , strain 013	196 – 265*
<i>Eimeria mitis</i> , strain 006	293 – 397*
<i>Eimeria praecox</i> , strain 007	293 – 397*
<i>Eimeria tenella</i> , strain 004	276 – 374*

\*Number of sporulated oocysts derived from precocious attenuated lines of coccidia, according to *in vitro* procedures of the manufacturer at the time of blending.

#### HIPRAMUNE T (solvent)

#### INDICATIONS:

For active immunisation of chicks from 1 day of age against coccidiosis caused by *Eimeria acervulina*, *Eimeria maxima*, *Eimeria mitis*, *Eimeria praecox* and *Eimeria tenella* to reduce typical signs of disease, intestinal lesions and oocysts output.

Onset of immunity: 14 days post-vaccination.

Duration of immunity: 63 days post-vaccination.

#### CONTRAINDICATIONS:

None.

#### ADVERSE REACTIONS:

None.

If you notice any side effects, even those not already listed in this package leaflet please inform your veterinary surgeon.

#### TARGET SPECIES:

Chickens.

#### DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION:

One dose of vaccine (0.007 ml) from 1 day of age.

Oral use.

The method of administration is by coarse spray.

#### ADVICE ON CORRECT ADMINISTRATION:

The method of administration is by coarse spray by using a suitable device (volume delivered: 28 ml/100 chicks, droplet size: 200–250 µm and working pressure: 2 to 3 bars). Before starting the preparation, make certain to have a clean container available with sufficient capacity for preparing the diluted vaccine suspension. Dilute the vaccine with the corresponding volumes:

DOSES	WATER	VACCINE	SOLVENT	TOTAL
1,000	223 ml	7 ml	50 ml	280 ml
5,000	1,115 ml	35 ml	250 ml	1,400 ml
10,000	2,230 ml	70 ml	500 ml	2,800 ml

Shake the solvent vial. Dilute the contents of the vial with clean room temperature water into an appropriate container.

Shake the vaccine vial and dilute the content into the previous solution.

Fill the reservoir of the spraying device with all the vaccine suspension prepared. Maintain the diluted vaccine suspension in continuous homogenisation by using a magnetic stirrer while the vaccine is being administered via coarse spray to the chicks.

To improve the uniformity of the vaccination maintain the chicks inside the transportation box for at least 1 hour in order to let them ingest all the vaccine droplets.

After this time, place the chicks carefully in the litter and continue with regular management practices.

#### WITHDRAWAL PERIOD:

Zero days.

#### SPECIAL STORAGE PRECAUTIONS:

Keep out of the sight and reach of children.

#### EVANT:

Store and transport refrigerated (2 °C - 8 °C). Do not freeze.

Shelf life of the veterinary medicinal product as packaged for sale: 10 months.

Shelf-life after dilution or reconstitution according to directions: 10 hours.

#### HIPRAMUNE T:

Store and transport refrigerated (2 °C - +8 °C). Do not freeze.

Shelf life of the veterinary medicinal product as packaged for sale: 24 months.

Do not use after the expiry date which is stated on the carton and the label.

#### SPECIAL WARNING(S):

##### Special warnings for each target species

The vaccine will not protect species other than chickens against coccidiosis and is only effective against the *Eimeria* species indicated.

##### Special precautions for use in animals

Chickens must be strictly floor-reared in the first 3 weeks after vaccination.

Vaccinate healthy chickens only.

It is recommended that litter should be removed and facilities and material cleaned between production cycles to reduce field infections.

##### Special precautions to be taken by the person administering the veterinary medicinal product to animals

Wash and disinfect hands and equipment after use.

##### Interaction with other medicinal products and other forms of interaction

No information is available on the safety and efficacy of this immunological veterinary medicinal product when used with any other veterinary medicinal product.

A decision to use this immunological veterinary medicinal product before or after any other veterinary medicinal product therefore needs to be made on a case by case basis.

No anticoccidial substances or other agents having anticoccidial activity via feed or water should be used for at least 3 weeks following vaccination of the chickens. The correct replication of the vaccine oocysts and consequently, the development of a solid immunity could be hindered. Additionally, the enhancement of protection produced by oocyst re-infections would also be limited.

##### Overdose (symptoms, emergency procedures, antidotes)

None.

##### Incompatibilities

Do not mix with any other veterinary medicinal product, except the solvent supplied for use with the veterinary medicinal product.

#### SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY:

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements.

#### OTHER INFORMATION:

##### Pack sizes:

Cardboard box with one vial of 1,000 doses (7 ml) and one vial with 50 ml of HIPRAMUNE T (solvent).

Cardboard box with one vial of 5,000 doses (35 ml) and one vial with 250 ml of HIPRAMUNE T (solvent).

Cardboard box with one vial of 10,000 doses (70 ml) and one vial with 500 ml of HIPRAMUNE T (solvent).

Not all pack sizes may be marketed.

Imported & Distributed by:

HIPRA PHILIPPINES, INC.

Unit 308 Venture Bldg.

corner Prime & Venture Street

Madrigal Business Park 2

Alabang, Muntinlupa City

VBPR No. R-2227  
Under veterinary prescription

FOR VETERINARY USE ONLY



LABORATORIOS HIPRA, S.A. Avda. la Selva, 135. 17170 Amer (Girona) Spain

Tel. (34) 972 43 06 60. Fax (34) 972 43 06 61. hipra@hipra.com