HIPRAVIAR S/H120

Live vaccine against Newcastle disease, La Sota strain, and Infectious bronchitis, H120 strain, in oral freeze-dried tablet

OUALITATIVE AND QUANTITATIVE COMPOSITION OF THE ACTIVES SUBSTANCES:

One dose (0.03 ml) contains:

Live Newcastle Disease Virus, strain La Sota ≥ 10^{6.5} EID₅₀ Infectious Avian Bronchitis Virus, strain H120 > 10³ EID₅₀

INDICATIONS FOR USE:

Poultry: To prevent Newcastle Disease and Infectious Bronchitis.

CONTRA-INDICATIONS:

None has been described.

ADVERSE REACTIONS:

Respiratory symptoms may occur in vaccinated birds at 3 days post-vaccination.

TARGET SPECIES:

Poultry.

DOSAGE FOR EACH SPECIES, ROUTE AND METHOD OF ADMINISTRATION:

Poultry: 1 dose/bird.

Ocular-nasal, oral or spray.

The veterinary surgeon will establish the most suitable vaccination programme according to the health condition of each farm and area.

ADVICE ON CORRECT ADMINISTRATION:

Ocular-nasal: Inject 10 ml of solvent to break the vacuum of the vial; before administration, shake gently until the freeze-dried tablet is complete resuspended. Mix the resuspended product with the rest of the solvent. Administer one drop of the vaccine (0.03 ml) per bird, in the eye or nostril, using a standard dropper (30 ml per 1,000 doses). **Oral:** Always open the vials under water, use a small quantity of water to reconstitute the freeze-dried tablet. Once the freeze-dried tablet is resuspended, pour it into an adequate container up to a volume of drinking water that can be ingested by the birds within 1 or 2 hours.

. Do not use water with residues or biocides, use fresh tap water only.

Spray: Validate the sprayer to check the amount of water needed. Always open the vials under water; use a small quantity of water to reconstitute the freeze-dried tablet. Once the freeze-dried tablet is resuspended, pour it into the amount of water needed.

- . Do not use water with residues of biocides, use fresh tap water only.
- . It is advisable to use coarse spray (larger than 150 microns) for primary vaccinations. For revaccionations, finer droplets can be used (smaller than 150 microns).
- . Spray administration requires the use of protective mask and glasses

WITHDRAWAL PERIOD:

0 days.

SPECIAL PRECAUTIONS FOR STORAGE:

Keep out of the reach and sight of children.

Store and transport refrigerated (+2 to +8 °C). Do not freeze. Protect from light.

SPECIAL WARNINGS:

Special warnings for each target species

Vaccinate only healthy animals.

If the number of birds does not coincide with the number of doses of available vials, an overdose should be administered, never administer less than the recommended dosage.

Special precautions for use in animals

Check the shelf life of the product.

Special precautions to be taken by the person administering the medicinal product to animals

Spray administration requires the use of protective mask and glasses.



LABORATORIOS HIPRA, S.A. Avda. la Selva, 135 . 17170 Amer (Girona) Spain

Tel. (34) 972 43 06 60. Fax (34) 972 43 06 61. hipra@hipra.com

713566-00.0 07-16

Use during pregnancy, lactation or lay

Can be administered in the laving period.

Interactions with other vaccines and medicinal products when administered in combination with the product

No information is available on the safety and efficacy of this vaccine when used with any other veterinary medicinal product. A decision to use this vaccine before or after any other veterinary medicinal product therefore needs to be made on a case by case basis.

Overdose

It causes no side effect.

Incompatibilities

In the absence of compatibility studies, do not mix with any other veterinary medicinal product.

SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED MEDICINAL PRODUCT OR WASTE MATERIALS DERIVED FROM THE USE OF SUCH PRODUCTS:

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with national requirements.

OTHER INFORMATION:

Pack sizes:

Pack with 10 vials of 1,000 doses. Pack with 10 vials of 2,500 doses. Not all pack sizes may be marketed.

wt5

Reg. n° 5/5/10/3667 Under veterinary prescription

FOR VETERINARY USE ONLY